

Flavonoids as nutraceuticals. A review

P. K. JAIN*, M. D. KHARYA, A. GAJBHIYE, U. V. S. SARA, V. K. SHARMA

D. J. College of Pharmacy
Modinagar, India

*corresponding author: phone: 09811767325,
e mail: pushpendra_utd@rediffmail.com, pushpendra.utd@gmail.com

Summary

Phenolic compounds form one of the main classes of secondary metabolites. They display a large range of structures and are responsible for the major organoleptic characteristics of plant-derived foods and beverages, particularly color and taste properties. They also contribute to the nutritional qualities of fruits and vegetables. Among these compounds, flavonoids constitute one of the most prevailing groups among plant phenolics. Owing to their importance in food organoleptic properties and human health, a better understanding of their structures and biological activities indicates their potentials as therapeutic agents and also for predicting and controlling food quality. Due to the variety of pharmacological activities in the mammalian body, flavonoids are more correctly referred as “nutraceuticals”.

Key words: bioflavonoids, structure-classification, nutraceuticals, antimicrobial activities, antioxidant activity, metabolic effects

INTRODUCTION

Phenolic compounds constitute one of the main classes of secondary metabolites. They display a large range of structures and they are responsible for the major organoleptic characteristics of plant-derived foods and beverages, particularly color and taste properties and they also contribute to the nutritional qualities of fruits and vegetables. The most important natural pigments are carotenoids which are tetrapyrrole derivatives of naturally occurring phenolic compounds ubiquitously distributed in plant kingdom. Among these compounds, flavonoids constitute one of the most ubiquitous groups of all plant phenolics. Up to now,

over 8,000 varieties of flavonoids have been identified [1]. Even 50 years ago, information on the working mechanisms of flavonoids was scarce. But it has been widely known for centuries that compounds of plant origin possess a broad spectrum of biological activity [2]. In 1930, Szent-Gyorgyi isolated a new substance from oranges and classified it as vitamin P, but later it became clear that this substance was actually a flavonoid [3]. Flavonoids drew greater attention of researchers with the discovery of the French Paradox, i.e., the decrease in cardiovascular incidents observed in the Mediterranean population which was associated to red wine consumption, and a greater amount of saturated fat in the average diet than in other countries [3].

STRUCTURE AND CLASSIFICATION OF FLAVONOIDS

Flavonoids occur as aglycones, glycosides and methylated derivatives. (4) In plants, flavonoids aglycones (i.e., flavonoids with no sugar attached) occur in a variety of structural forms. All of them contain fifteen carbon atoms in their basic nucleus: two six-membered rings linked with a three carbon unit which may or may not be a part of a third ring [5]. For convenience, the rings are labeled A, B, and C (fig. 1). The individual carbon atoms are based on a numbering system which uses ordinary numerals for the A and C and “primed” numerals for B-ring [1]. Primed modified numbering system is not used for chalcones [2] and the isoflavones derivatives [6]; the pterocarpan and the rotenoids [6]. The different ways to close this ring associated with the different oxidation degrees of ring A provide the various classes of flavonoids. The six-membered ring condensed with the benzene ring is either a pyrone flavones [1] flavonols [3] or its dihydroderivative flavanones (4) and flavan-3-ols (5). The position of the benzenoid substituent divides the flavonoids into two classes: flavonoids [1] (2nd position) and isoflavonoids [6] (3rd position). Most of flavonoids occur naturally associated with sugar in conjugated form and, within any one class, may be characterized as monoglycosidic, diglycosidic, etc. The glycosidic linkage is normally located at position 3 or 7 and the carbohydrate unit can be L-rhamnose, D-glucose, glucorhamnose, galactose or arabinose [8].

FLAVONOIDS AS NUTRACEUTICALS

“Nutraceutical” is a term coined in 1979 by Stephen DeFelice. (9) It is defined “as a food or parts of food that provide medical or health benefits, including the prevention and treatment of disease.” Nutraceuticals may range from isolated nutrients, dietary supplements, and diets to genetically engineered “designer” food, herbal products and processed products such as cereals, soups, and beverages. A nutraceutical is any nontoxic food extract supplement that has scientifically proven health benefits for both the treatment and prevention of disease [10]. The increasing interest in nutraceuticals reflects the fact that consumers hear about

epidemiological studies indicating that a specific diet or component of the diet is associated with a lower risk for a certain disease. The major active nutraceutical ingredients in plants are flavonoids. As is typical for phenolic compounds, they can act as potent antioxidants and metal chelators. They also have long been recognized to possess anti-inflammatory, antiallergic, hepatoprotective, antithrombotic, antiviral, and anticarcinogenic activities, as discussed in the subsections that follow (fig. 1, fig. 2, tab. 1):




Figure 1. Chemical structures of some representative flavonoids



Figure 2. General structures for various classes of flavonoids

Table 1.

Substitution patterns of series of flavonoids									
Group	3	5	6	7	8	3'	4'	5'	C ₂ =C ₃
Flavones									
Apigenin	H	OH	H	OH	H	H	OH	H	+
Diosmin	H	OH	H	Oru	H	OH	OH	H	+
Luteolin	H	OH	H	OH	H	OH	OH	H	+
Flavonol									
Quercetin	OH	OH	H	OH	H	OH	OH	H	+
Kaempferol	OH	OH	H	OH	H	H	OH	H	+
Galangin	OH	OH	H	OH	H	H	H	H	+
Fisetin	OH	H	H	OH	H	OH	OH	H	+
Myricetin	OH	OH	H	OH	H	OH	OH	OH	+
Vitexicarpin	OCH ₃	OH	OCH	OCH ₃	H	OH	OCH ₃	H	+
Flavanone									
Naringenin	H	OH	H	OH	H	H	OH	H	-
Eriodictyol	H	OH	OH	OH	H	OH	OH	H	-
Pinocembrin	H	OH	H	OH	H	H	H	H	-
Liquiritigenin	H	H	H	OH	H	H	OH	H	-
Flavanonol									
Taxifolin	H	OH	H	OH	H	OH	OH	H	-
Isoflavone									
Genistein	-	OH	H	OH	H	H	OH	H	+
Tectorigenin	-	OH	OCH ₃	OH	H	H	OH	H	+
Daidzein	-	H	H	OH	H	H	OH	H	+
Formononetin	-	H	H	OH	H	H	OCH ₃	H	+
Flavan-3-ols									
(+) Catechin	βOH	OH	H	OH	H	OH	OH	H	-
(-) Epicatechin	αOH	OH	H	OH	H	OH	OH	H	-
(-) Epigallocatechin	αOH	OH	H	OH	H	OH	OH	OH	-
Flavylium Salts									
Cyanidin	OH	OH	H	OH	H	OH	OH	H	-
Pelargonidine	OH	OH	H	OH	H	H	OH	H	-

Antioxidant activity (tab. 2, tab. 3)

The best-described property of almost every group of flavonoids is their capacity to act as antioxidants. The flavones and catechins seem to be the most powerful flavonoids protecting the body against reactive oxygen species (ROS). Body cells and tissues are continuously threatened by the damage caused by free radicals and ROS which are produced during normal oxygen metabolism or are induced by exogenous damage [11, 12]. Free radicals and ROS have been implicated in a large number of human diseases [13, 14]. Quercetin, kaempferol, morin, myricetin and rutin, by acting as antioxidants, exhibited beneficial effects such as anti-inflammatory, antiallergic, antiviral as well as anticancer activity. They have also been suggested to play a protective role in liver diseases, cataracts and cardiovascular diseases. Quercetin and silybin, acting as free radical scavengers, were

shown to exert a protective effect in liver reperfusion ischemic tissue damage [15, 16]. The scavenging activity of flavonoids has been reported to be in the order: myricetin > quercetin > rhamnetin > morin > diosmetin > naringenin > apigenin > catechin > 5,7-dihydroxy-3',4',5'-trimethoxyflavone > robinin > kaempferol > flavone [17].

Table 2.

Reactive oxygen species that can be scavenged or whose formation can be inhibited by flavonoids [18, 19]

O_2^- (Superoxide anion)	One-electron reduction product of O_2 . Produced by phagocytes, formed in autoxidation reactions (flavoproteins, redox cycling), and generated by oxidases (heme proteins).
HO_2^-	Protonated form of O_2^- .
H_2O_2 (Hydrogen Peroxide)	Two-electron reduction product of O_2 formed from O_2 by O_2^- dismutation or directly from O_2 . Reactivity of O_2^- and H_2O_2 is amplified in the presence of heme proteins.
OH^\cdot (Hydroxy radical)	Three-electrons reduction product of O_2 generated by Fenton reaction, transition metal (iron, copper)-catalysed Haber-Weiss reaction; also formed by decomposition of peroxyinitrite produced by the reaction of O_2^- with NO^\cdot (Nitric oxide radical). Example: Lipid radical (LO^\cdot).
RO^\cdot (Alkoxy radical) ROO^\cdot (Peroxy radical)	Example: Lipid peroxy radical (LOO^\cdot) produced from organic hydroperoxide (e.g. lipid hydroperoxide, $LOOH$), $ROOH$ by hydrogen abstraction.
1O_2	Singlet oxygen

Table 3:

Characteristics of flavonoid structure for most effective radical-scavenging activity [20-22]

1. The catechol (O-dihydroxy) group in the ring confers great scavenging ability.
2. A pyrogallol (trihydroxy) group in ring B of a catechol, as in myricetin, produces even higher activity. The C2-C3 double bond of the C ring appears to increase scavenger activity because it confers stability to the phenoxyl radical produced.
3. The 4-oxo (keto double bond at position 4 of the C ring), especially in association with the C2-C3 double bond, increases scavenger activity by delocalizing electrons from Bring.
4. The 3-OH group on the C ring generates an extremely active scavenger; in fact, the combination of C2-C3 double bond and 4-oxo group appears to be the best combination on the top of the catechol group.
5. The 5-OH and 7-OH groups may also add scavenging potential in certain cases.

Antimicrobial activity

Flavonoids and esters of phenolic acids have also been investigated for their antibacterial, antifungal and antiviral activities.

Antibacterial activity

Antibacterial activity has been displayed by a number of flavonoids. Quercetin has been reported to completely inhibit the growth of *Staphylococcus aureus*. Most of the flavanones having no sugar moiety showed antimicrobial activities whereas none of the flavonols and flavonolignans tested showed inhibitory activity on microorganisms [23].

Antifungal activity

A number of flavonoids isolated from the peelings of tangerine orange, when tested for fungistatic activity towards *Deuterophoma tracheiphila* were found to be active; nobiletin and langeritin exhibited strong and weak activities, respectively, while hesperidin could slightly stimulate fungal growth. Chlorflavonin was the first chlorine-containing flavonoid-type antifungal antibiotic produced by strains of *Aspergillus candidus* [24].

Antiviral activity

Naturally occurring flavonoids with antiviral activity have been recognized since the 1940s but only recently attempts have been made to make synthetic modifications of natural compounds to improve antiviral activity. Quercetin, morin, rutin, dihydroquercetin (taxifolin), apigenin, catechin and hesperidine have been reported to possess antiviral activity against some of 11 types of viruses [25]. The antiviral activity appears to be associated with the nonglycosidic compounds and hydroxylation at the 3-position is apparently a prerequisite for antiviral activity. It has been found that flavonols are more active than flavones against *Herpes simplex* virus type 1. The order of importance was galangin > kaempferol > quercetin [26]. Recently, a natural plant flavonoid polymer of molecular weight 2,100 daltons was found to have antiviral activity against two strains of type 1 Herpes simplex virus and type 2 Herpes simplex viruses²⁷. Because of the worldwide spread of HIV since the 1980s, the investigation of the antiviral activity of flavonoids has mainly focused on HIV [28]. There have appeared several recent reports on the anti-AIDS activity of flavonoids. Out of twenty eight flavonoids tested, the flavans were generally more effective than flavones and flavanones in the selective inhibition of HIV-1 and HIV-2 or similar immunodeficiency virus infections [29].

Effect on gastrointestinal system

Antiulcer activity exerts significant anti-inflammatory activity in the animal model of both acute and chronic inflammation when given orally or topically [34, 35]. Hesperidin, a citrus flavonoid, possesses significant anti-inflammatory and analge-

sis effects [36]. Recently, apigenin, luteolin and quercetin have been reported to exhibit anti-inflammatory activity [37]. A number of reports have been published which demonstrate that flavonoids can modulate arachidonic acid metabolism via the inhibition of cyclo-oxygenase (COX) and lipoxygenase activity (LO). It has been also speculated that the anti-inflammatory and antiallergic properties of flavonoids are the consequence of their inhibitory actions on arachidonic acid metabolism [38]. Some recent studies have indicated that flavonoids possess antiulcerogenic activity. Flavonoid glycosides of *Ocimum basilicum* (Labiatae) decreased ulcer index and inhibited gastric acid and pepsin secretions in aspirin-induced ulcers in rats [30]. Quercetin, rutin, and kaempferol administered intraperitoneally (25–100 mg/kg) inhibited dose-dependent gastric damage produced by acidified ethanol in rats [31].

Hepatoprotective activity

The liver is subject to acute and potentially lethal injury by several substances including phalloidin (the toxic constituent of the mushroom, *Amanita phalloides*), CCl_4 , galactosamine, ethanol and other compounds. Flavonoids have also been found to possess hepatoprotective activity. In the study carried out in order to investigate the flavonoid derivatives: silymarin, apigenin, quercetin, and naringenin, as putative therapeutic agents against microcrystin LR-induced hepatotoxicity, silymarin was found to be the most effective [32]. The flavonoid, rutin and venoruton showed regenerative and hepatoprotective effects in experimental cirrhosis [33].

Antiinflammatory activity

The antiinflammatory activity of flavonoids has been reported in many animal models. Flavone/flavonol glycosides as well as flavonoid aglycones have been reported to possess LO and COX inhibitory activities [39, 40].

Antidiabetic effects

Flavonoids, especially quercetin, have been reported to possess antidiabetic activity. Vessal et al reported that quercetin brings about the regeneration of pancreatic islets and probably increases insulin release in streptozotocin induced diabetic rats [41]. Also in another study, Hif and Howell reported that quercetin stimulates insulin release and enhanced Ca^{2+} uptake from isolated islets cell which suggests a place for flavonoids in noninsulin-dependent diabetes [42, 43].

Effect of vasorelaxant agent on cardiovascular system

The consumption of flavonoids may prevent endothelial dysfunction by enhancing the vasorelaxant process leading to the reduction of arterial pressure [44, 45]. Endothelial dysfunction represents a critical event in the development of cardiovascular diseases and the major complication of atherosclerosis and arterial thrombus formation [46].

The consumption of flavonoids can prevent a number of cardiovascular diseases including hypertension and atherosclerosis [47, 48]. Recently, many experimental studies have shown that these polyphenolic compounds may reduce the arterial pressure in rats and enhance the vasorelaxant process. The endothelium-dependent relaxation induced by flavonoids has been well documented. Furthermore, investigators have demonstrated that *Anthocyanin delphinidin* exerts a significant endothelium dependent vasorelaxation [49, 50].

Antiatherosclerotic effects

Oxidative modification of low-density lipoproteins (LDL) by free radicals is an early event in the pathogenesis of atherosclerosis effect [52]. Furthermore, a Japanese study reported an inverse correlation between flavonoid intake and total plasma cholesterol concentrations [53].

Antithrombogenic effects

Platelet aggregation plays a pivotal role in the physiology of thrombotic diseases. Activated platelets adhering to vascular endothelium generate lipid peroxides and oxygen free radicals which inhibit the endothelial formation of prostacyclin and nitrous oxide. It was shown in the 1960s that tea pigment can reduce blood coagulability, increase fibrinolysis and prevent platelet adhesion and aggregation [54]. Selected flavonoids, such as quercetin, kaempferol and myricetin were shown to be effective inhibitors of platelet aggregation in dogs and monkeys [55]. Flavonols are particularly antithrombotic because they directly scavenge free radicals, thereby maintaining proper concentration of endothelial prostacyclin and nitric oxide [56]. One study showed that flavonoids are powerful antithrombotic agents *in vitro* and *in vivo* because of their inhibition of the activity of cyclooxygenase and lipoxigenase pathways [57].

Cardioprotective effects

Recently, the interest in flavonoids has been stimulated by the potential health benefits arising from the antioxidant activity of these polyphenolic compounds.

These are the results of their high propensity to transfer electrons, chelate ferrous ions and scavenge reactive oxygen species [58]. Due to these properties, flavonoids have been considered as potential protectors against chronic cardiotoxicity caused by the cytostatic drug – doxorubicin. Doxorubicin is a very effective anti-tumor agent, although, its clinical use is limited by the occurrence of a cumulative dose-related cardiotoxicity, resulting in, for example, congestive heart failure (negative inotropic effect). In the recent report, the cardiotoxicity of doxorubicin on the mouse left atrium has been inhibited by flavonoids, 7- monohydroxyethylrutoside and 7', 3', 4'- trihydroxyethylrutoside [34, 59-61].

Antineoplastic activity

A sufficient number of flavonoids have exhibited antineoplastic activity. Several recent reviews have highlighted this activity. Detailed studies [62-64] have revealed that quercetin exerted a dose-dependent inhibition of growth and colony formation. The flavonoids: kaempferol, catechin, toxifolin and fisetin, also suppressed cell growth [65, 66]. On screening the antileukaemic efficacy of 28 naturally occurring and synthetic flavonoids on human promyelocytic leukaemic HL-60 cells, genistein, an isoflavone was found to have strong effect [67, 68].

Effect on central nervous system

Synthetic flavonoids, such as 6-bromoflavone and 6-bromo-3'-nitroflavones, were shown to displace [3H] flumazenil binding to membranes from rat cerebellum but not from spinal cord, indicating selectivity for the BZOmega receptor subtype, but the latter was more potent than 6-bromoflavone. Results from two conflict tests in rats showed that these synthetic flavonoids possess anxiolytic like properties similar or superior to that of diazepam [69].

Toxicity of flavonoids

Flavonoids are ubiquitous in plant foods and drinks and, therefore, a significant quantity is consumed in our daily diet. The toxicity of flavonoids is very low in animals. For rats, the LD50 is 2–10 g per animal for most flavonoids. Similar doses in humans are quite unrealistic. As a precaution, doses less than 1mg per adult per day have been recommended for humans [70]. Dunnick and Hailey reported that high doses of quercetin over several years might result in the formation of tumors in mice [71]. However, in other long-term studies, no carcinogenicity was found [72]. Moreover, as described earlier, quercetin has been reported to be anti-mutagenic *in vivo*.

CONCLUSION

Flavonoids comprise a vast array of biologically active compounds that are ubiquitous in plants, many of which have been used in traditional eastern medicine for thousands of years. They also constitute an unavoidable component of the diet. In the present review, we have reviewed detailed structural aspects and biological properties of flavonoids. The chemical and structural similarities of flavonoids with numerous biomolecules as well as their crucial role in plant-insect and plant-bacterial interactions make them an attractive class of phytoconstituents for biological activity. Their widespread occurrence, broad spectrum diversity and natural origin make them appropriate chemical scaffolds for novel therapeutic agents. Of the many actions of flavonoids, antioxidant and antiproliferative effects stand out. Given that certain substituents are known to be required or increase their actions, the therapeutic potential of selected flavonoids is fairly obvious. These natural compounds have several great advantages over other therapeutic agents for the following reasons:

1. Many diets are rich in these phenolics and are daily consumed.
2. They rarely have any side effects.
3. They have relatively long half-life.
4. They can be easily absorbed in the intestine after ingestion.

The study of flavonoids is complex because of the heterogeneity of different molecular structures and the scarcity of data on bioavailability. There is a need to improve analytic techniques to allow collection of more data on absorption and excretion. Data on the long-term consequences of chronic flavonoid ingestion are especially scarce. Finally, we think that natural, hemisynthetic and synthetic flavonoids alone or in combination with other preventive and/or therapeutic strategies will become effective future drugs against the most common degenerative diseases such as cancer, diabetes and cardiovascular complications.

REFERENCES

1. De Groot H, Raven U. Tissue injury by reactive oxygen species and the protective effects of flavonoids. *Fundam Clin Pharma Col* 1998; 12:249-55.
2. Robak J, Gryglewski RJ. Bioactivity of flavonoids. *Pol J Pharmacol* 1996; 48:555-64.
3. Ranaud S, de Lorgeril M. Wine, alcohol, platelets, and the french paradox for coronary heart disease. *Lancet* 1992; 339(8808):1523-6.
4. Harborne JB. *The flavonoids – Advances in Research since 1980*. Ed 1. London 1988.
5. Middleton E. The flavonoids. *Trends Pharmacol Sci* 1984; 5:335-8.
6. Havsteen B. Flavonoids, a class of natural products of high pharmacological potency. *Biochem Pharmacol* 1983; 32(7):1141-8.
7. Harborne JB. Nature, distribution and function of plant flavonoids. In: Cody V, Middleton Elliott Jr, Harborne JB (eds.). *Plant flavonoids in biology and medicine: biochemical, pharmacological and structure activity relationship*. New York 1986:15-24.
8. Harborne JB, Baxter, H. *The handbook of natural flavonoids*. Vol.1-2. New York 1999.
9. DeFelice SL. Nutraceuticals: Opportunities in an Emerging Market. *Scrip Mag* 1992; 9:xxx-yyy.

10. Dillard, CJ, German JB. Phytochemicals: nutraceuticals and human health. *J Sci Food Agric* 2000; 80:1744-56.
11. De Groot H. Reactive oxygen species in tissue injury. *Hepatogastroenterology* 1994; 41:328-32.
12. Grace PA. Ischaemia-reperfusion injury. *Br J Surg* 1994; 81:637-47.
13. Wegener T, Fintelmann V. Flavonoids and bioactivity. *Wein Med Wochem Schr* 1999; 149:241-7.
14. Ares JJ, Outt PE. Gastroprotective agents for the prevention of NSAID induced gastropathy. *Curr Pharm Des* 1998; 4:7-36.
15. Hillwell B. Free radicals, antioxidants and human disease: Curiosity, cause or constipation. *Lancet* 1994; 344:721-4.
16. Fraga CG, Mactino US, Ferraro GE, Coussio JF, Boveris A. Flavonoids as antioxidants evaluated by *in vitro* and *in situ* liver chemiluminescence. *Biochem Med Metabol Biol* 1987; 36:717-20.
17. Ratty AK. Effects of flavonoids on nonenzymatic lipid peroxidation: structure activity relationship. *Biochem Med Metabol Biol* 1988; 39:67-79.
18. Bors W, Heller W, Michel C, Saran M. Flavonoids as antioxidant: Determination of radical scavenging efficiencies. *Methods Ezymol*, 1990; 186:343-355.
19. Cotellet N, Bernier JL, Catteau JP, Pommery J, Wallet JC, Gaydou EM. Antioxidant activity of hydroxy flavonoids. *Free Radic Biol Med* 1996; 20:35-43.
20. Rice-Evans CA, Nicholas JM, Paganga G. Structure antioxidant activity relationships of flavonoids and phenolic acid. *Free Radic Biol Med* 1996; 20(7):933-56.
21. Amic D, Davidovic-Amic D, Beslo D, Trinajstic N. Structure radical scavenging activity relationships of flavonoids. *Croat Chem Act* 2003; 76:55-61.
22. Farkas O, Jakus J, Heberger K. Quantitative structure-antioxidant activity relationship of flavonoid compounds. *Molecules* 2004; 9:1079-88.
23. Havsteen B. Flavonoids: A class of natural products of high pharmacological potency. *Biochem Pharmacol* 1983; 32(7):1141-8.
24. Tencate JW, van Hoeringen NJ, Gerritsen J, Glasius E. Biological activity of a semisynthetic flavonoid O-(hydroxyethyl) rutosine: Light scattering and metabolic studies of human red cells and platelets. *Clin Chem* 1973; 19:31-5.
25. Selway JWT. Antiviral activity of flavones and flavans. In: Cody V, Middleton E, Harborne JB (eds.). *Plant flavonoids in biology and medicine: Biochemical, pharmacological and structure activity relationships*. New York 1986; 521-36.
26. Thomas PRS, Nash GB, Dormandly JA. White cells accumulation in dependent legs of patients with venous hypertension: A possible mechanism for trophic changes in the skin. *Br Med J* 1988; 296:1673-95.
27. Loewenstein WR. Junctional Intercellular communication and the control of growth. *Biochem Biophys Acta* 1979; 560:1-65.
28. Ng TB, Huang B, Fong WP, Yeung HW. Anti-human immunodeficiency virus (anti-HIV) natural products with special emphasis on HIV transcriptase inhibitors. *Life Sci* 1997; 61:933-49.
29. Gerdin B, Srenso E. Inhibitory effect of flavonoids on increased microvascular permeability induced by various agents in rat skin. *Int J Micro Cir Clin Exp* 1983; 2:39-46.
30. Alarcon DL, Martin MJ, Locasa C, Motilva V. Antitumorogenic activity of flavonoids and gastric protection. *Ethnopharmacol* 1994; 42:161-70.
31. Izzo AA, Carlo GD, Mascolo N, Capasso F, Autore G. Effect of quercetin on gastrointestinal tract. *Phyto Ther Res* 1994; 8:179-85.
32. Carlo GD, Autore G, Izzoaa et al. Inhibition of intestinal motility and secretion by flavonoids in mice and rats; structure activity relationships. *J Pharm Pharmacol* 1993; 45:1045-59.
33. Lorenz W, Kusche J, Barth H, Mathias CH. Action of several flavonoids on enzyme of histidine metabolism *in vivo*. In: Cz. Maslinski (ed.). *Histamine*. Pennsylvania 1994; 265-9.
34. Lee SJ, Son KH, Chang HW et al. Anti-inflammatory activity of naturally occurring flavone and flavonol glycosides. *Arch Pharm Res* 1993; 16(1):25-8.
35. Hang T, Jin JB, Cho S, Cyang JC. Evaluation of anti-inflammatory effects of baicalein on dextran sulfate sodium-induced colitis in mice. *Planta Med* 2002; 68(2):268-71.
36. Shahid F, Yang Z, Saleemi ZO. Natural flavonoids as stabilizers. *J Food Lipids* 1998; volume:69-75.
37. Farmica JV, Regelson W. Review of the biology of quercetin and related bioflavonoids. *Fd Chem Toxic* 1995; 33(12):1061-80.

38. Ferrandiz ML, Alcaraz MJ. Anti-inflammatory activity and inhibition of arachidonic acid metabolism by flavonoids. *Agent and Actions* 1991; 32(3/4):283-8.
39. Kim HP, Mani I, Iversen L, Ziboh VA. Effects of naturally occurring flavonoids and bioflavonoids on epidermal cyclooxygenase and lipoxygenase from guinea pigs. *Prostag Leukot Essent Fatty Acids* 1998; 58:17-24.
40. Jachak SM. Natural products: Potential source of COX inhibitors. *CRIPS* 2001; 2(1):12-15.
41. Vessal M, Hemmati M, Vasei M. Antidiabetic effects of quercetin in streptozocin induced diabetic rats. *Comp Biochem Physiol* 2003; 135:357-64.
42. Hif CS, Howell SL. Effects of epicatechin on rat islets of langerhans. *Diabetes*, 1984; 33:291-296.
43. Hif CS, Howell SL. Effects of flavonoids on insulin secretion and $45\text{Ca}+2$ handling in rat islets of langerhans. *J Endocrinol* 1985; 107:1-8.
44. Iijima K, Aviram M. Flavonoids protect LDL from oxidation and attenuate atherosclerosis. *Curr Opin Lipidol* 2001; 12:41-48.
45. Bernatova I, Pechanova O, Balal P et al. Wine polyphenols improve cardiovascular remodeling and vascular function in NO-deficient hypertension. *Am J Physiol Heart Cir Physiol* 2002; 282:942-8.
46. Jayakody TL, Senaratne MPJ, Thompson ABR, Kappagoda CT. Cholesterol feeding impairs endothelium-dependent relaxation of rabbit aorta. *Can J Physiol Pharmacol* 1985; 63:1206-9.
47. Hertog MG, Feskens EJ, Hallman PC, Katan MB, Kromhout D. Dietary antioxidant flavonoids and risk of coronary heart disease: the Zutphen elderly study. *Lancet* 1993; 342:1007-11.
48. Hertog MG, Hallman PC, Katan MB, Kromhout D. Intake of potentially anticarcinogenic flavonoids and their determinants in adults in the Netherlands. *Nutr Cancer* 1993; 20:21-9.
49. Andriambeloson E, Kleschyov AI, Muller B et al. Nitric oxide production and endothelium dependent vasorelaxation induced by wine polyphenols in rat aorta. *Br J Pharmacol* 1997;120:1053-8.
50. Burns J, Gardner PT, O'Neil J et al. Relationship among antioxidant activity, vasodilation capacity and phenolic content of red wines. *J Agric Food Chem* 2000; 48:220-30.
51. De-Whalley C, Rankin SM, Houct JRS et al. Flavonoids inhibit the oxidative modification of low density lipoprotein by macrophages. *Biochem Pharmacol* 1990; 39:1743-50.
52. Fuhrman B, Lavy A, Aviram M. Consumption of red wine with meals reduces the susceptibility of human plasma and low-density lipoproteins to lipid peroxidation. *Am Soc Nutr* 1995; 61:549-54.
53. Arai Y, Watanabe S, Kimira M et al. Dietary intake of flavonols, flavones and isoflavones by Japanese women and the inverse correlation between quercetin intake and plasma LDL cholesterol concentration. *J Nutr* 2000; 130:2243-50.
54. Lou FQ, Zhang MF, Zhang XG, Liu JM, Yuan WL. A study on tea pigment in prevention of atherosclerosis. *Chin Med J (Engl)* 1989; 102:579-83.
55. Osman HE, Maalej N, Shanmuganayagam D, Folts JD. Grape juice but not orange or grapefruit juice inhibits plate activity in dogs and monkey's. *J Nutr* 1998; 128:2307-12.
56. Gryglewski RJ, Korbut R, Robak J, Swips J. On the mechanism of anti thrombotic action of flavonoids. *Biochem Pharmacol* 1987; 36:317-22.
57. Alcaraz MJ, Ferrandiz ML. Modification of arachidonic metabolism by flavonoids. *J Ethnopharmacol* 1987; 21:209-29.
58. Kandaswami C, Middleton E. Free radical scavenging and antioxidant activity of plant flavonoids. In: Armstrong D ed. *Free radicals in diagnostic medicine*, New York: Plenum Press, Ed 2, 1994. 351-376.
59. Lackeman GM, Claeys M, Rwanagabo PC, Herman AG, Vlietinck A. Chronotropic effect of quercetin on guinea pig right atrium. *J Planta Med* 1986; 52:433-9.
60. Huesken BCP, Dejong J, Beekman B, Onderwater RCA. Flavonoids as cardioprotective agents. *Cancer Chemotheapy Pharmacol* 1995; 37:55-62.
61. Bast A, Kaiserov H, den Hartog GJM, Haenen GRMM, van der Vijgh WJF. Protectors against doxorubicin-induced cardiotoxicity: Flavonoids. *Cell Biol Toxicol* 2007; 23:39-47.
62. Kontruck SJ, Radecki T, Brozowski T et al. Antiulcer and gastroprotective effects of solon, a synthetic flavonoid derivative of sophorandiol. Role of endogenous prostaglandins. *Bur J Pharmac* 1986; 125:185-92.
63. Izzo AA, Dicarlo, G, mascolo N, Capasso F, Autore G. Antiulcer effects of flavonoids. Role of endogenous PAF. *Phytotherapy Res* 1991; 8:179-81.
64. Murakami S, Muramatsu M, Otomo S. Gastric H⁺/K⁺ ATPase inhibition by catechins. *J Pharm Pharmacol* 1992; 44:926-8.

65. Kim HK, Namgoong SY, Kim HP. Biological actions of flavonoids-I. Arch Pharmacol Res 1993; 16:18-27.
66. Gill B, Sanz MJ, Terencio MC, Ferrandiz ML, Bustos G, Paya M. The flavonoids. Life Sci 1994; 54:333-9.
67. Hirano T, Gotoh M, Oak K. Natural flavonoids and lignans are plant cytostatic agents against Human Leukemic HL-60 cells. Life Sci 1994; 55:1061-9.
68. Wei H, Tye L, Bresnick E, Birt DF. Inhibitory effect of Apigenin, a plant flavonoids on epidermal ornithine decarboxylase skin tumor promotion in mice. Cancer Res 1990; 50: 499-502.
69. Griebel G, Perrault G, Tan S, Schoemaker H, Sanger DJ. Pharmacological studies on synthetic flavonoids: Comparison with diazepam. Neuropharmacol 1999; 38:965-77.
70. Starvic B. Mutagenic food flavonoids. Fed Proc 1984; 43:2344.
71. Dunnick JK, Hailey JR. Toxicity and carcinogenicity studies of quercetin, a natural component of food. Fundam Appl Toxicol 1992; 19:423-31.
72. Plakas SM, Lee TC, Wolke RE. Absence of overt toxicity from feeding the flavonol, quercetin rainbow trout (*Calmo gairdneri*). Food Chem Toxicol 1985; 23:1077-80.

FLAWONOIDY JAKO NUTRACEUTYKI – PRZEGLĄD

P. K. JAIN*, M. D. KHARYA, A. GAJBHIYE, U. V. S. SARA, V. K. SHARMA

D. J. College of Pharmacy
Modinagar, India

*autor, do którego należy kierować korespondencję : tel.: 09811767325,
e mail: pushpendra_utd@rediffmail.com, pushpendra.utd@gmail.com

Streszczenie

Związki fenolowe są jedną z głównych grup metabolitów wtórnych. Mogą mieć bardzo różną strukturę i są odpowiedzialne za większość cech organoleptycznych żywności pochodzenia roślinnego i napojów, szczególnie za ich właściwości kolorystyczne i smakowe. Podwyższają także wartości odżywcze owoców i warzyw. Flawonoidy są jedną z największych grup fenoli roślinnych. Lepsze zrozumienie budowy i właściwości biologicznych tych związków może wykazać ich potencjał jako związków leczniczych, a także mających znaczenie w jakości żywności. Ze względu na różnorodność działań farmakologicznych w organizmach ssaków, flawonoidy powinno się nazywać „nutraceutykami”.

Słowa kluczowe: bioflawonoidy, klasyfikacja ze względu na budowę, nutraceutyki, działanie przeciwbakteryjne, działanie przeciwutleniające, wpływ na metabolizm

