

Microbial transformation of α -naphthol by *Aspergillus niger* – PTCC 5011

AKBAR ESMAEILI^{1*}, SHAHROUZ FAZELI¹, NASRIN MOAZAMI²

¹Department of Chemical Engineering,
North Tehran Branch
Islamic Azad University
P.O. Box 19585-936
Tehran, Iran

²Department of Biotechnology
Iranian Research Organization for Science and Technology
Teheran, Iran

S u m m a r y

The metabolism of α -naphthol by *Aspergillus niger* PTCC 5011, a fungi isolated in Ardebil, Iran, from industrial wastewater, was studied. *A. niger* metabolized approximately 80% of α -naphthol within 5 days. The identification and quantification of degradation products using GC-MS demonstrated that approximately 41% of the parent compound was converted into 1-ethyl-2-methyl benzene, 7.43% was converted into acetoneaphthone, 5.55% was transformed into 4-hydroxy-1-naphthyl sulfate, 3% into 1,4-naphthoquinone, and about 6.68% into 2-phenyl-1,2,3-tetrahydro-1-naphthol. These results support a role for *A. niger* in affecting the environmental fat of pollutants in ecosystems.

Key words: biotransformation, bioconversion, *Aspergillus niger*, α -naphthol, fungi

INTRODUCTION

The polycyclic aromatic hydrocarbon naphthalene is used in the production of phthalate plasticizers and resins, azo dyes, dispersants, and tanning agents in the rubber and leather industries [1]. Naphthol is a common bicyclic aromatic often released into the environment [2]. Naphthol is often liberated into the environment as a result of the oxidation of naphthalene by certain fungi and bacteria [2, 5, 6] and is particularly important both as a synthetic precursor of the

insecticide Sevin (1-naphthyl-N-methylcarbamate) and as a degradation product of this compound via chemical and biological processes [2, 5, 6, 8]. It has toxic effects in aquatic ecosystems, especially on marine invertebrates [15].

Due to its toxicity to marine life [8] and human beings [18, 21], industrial wastewater containing α -naphthol must be treated before it is discharged into or reused in the environment [16]. Fungi play an important role in the metabolism of many chemicals, including aromatic hydrocarbons, in both aquatic and terrestrial environments [7, 14, 19, 21]. There are studies attesting to the ability of strains of different bacteria, e.g., *Brevibacterium* spp. HL 4, *Pseudomonas* spp. DLC-P11, *Arthrobacter sulphureus* RKJ4, *Acidovorax delafieldii* P4-1 [17], *Halosphaeria mediose-tigera*, *Culcitalna achraspora*, *Humicola alopallonella*, *Brevibacterium* spp., *Flavobacterium* spp., *Serratia marina*, *Spirillum* spp., *Candida parapsilosis*, *Rhodotorula glutinis*, *Trichosporon ferments*, *Aspergillus fumigatus* [22], *Micrococcus* spp. [18], *Heliscus lugdunensis* [7], and *Rhodococcus* spp. [2], to disintegrate α -naphthol and reports about isolation and examination of these fungal strains with respect to their α -naphthol tolerance. However, we are not aware of studies describing biotransformation of α -naphthol by *A. niger* PTCC 5011. This paper describes the ability of *A. niger* PTCC 5011 to transform this compound and the identification of pathways involved in the biotransformation of the resultant metabolites. In a previous study, microbial transformation of some monoterpenes (citral, geraniol, nerol, menthol, and myrcene) by various fungi, including *Aspergillus niger*, *Pseudomonas* sp., and *Penicillium* sp., produced a number of compounds, such as 1,8-cineole, 2,6-dimethyl-octane, α -pinene, α -terpineol, cis-p-menthan-7-ol, dihydrolinalool, γ -terpinene linalool, limonene, *p*-cymene, *p*-menth-1-ene, sabinene, and *trans-p*-menthan-1-ol [24-31].

MATERIALS AND METHODS

Chemicals

All chemicals were of analytical grade (gradient grade in the case of chromatography solvents). α -Naphthol was purchased from Panreac Química S.A.U. (PS) of Barcelona, Spain (>99% purity). Tween 80 was provided by the Department of Chemistry, North Tehran Branch, Islamic Azad University, Tehran, Iran. Czapek-Dox broth (Czapek's medium) was purchased from Difco (Detroit, Michigan, USA). All other chemicals were purchased from Merck (Darmstadt, Germany).

Microorganism

A strain of *Aspergillus niger* was isolated from the wastewater of an industrial plant in Ardebil, and was identified according to its physiological and morphological

characteristics. *Aspergillus niger* (PTCC 5011) was identified according to the Persian Type Culture Collection, Iranian Research Organization for Science & Technology, Tehran, Iran. The fungus mycelia was stored on agar slants at 4°C.

Culture condition

Spores and mycelia (2×10^8 conidia/ml medium) were aseptically inoculated into 250 ml Erlenmeyer flasks containing 100 ml of Czapek-Dox broth. The Czapek's medium (in g L⁻¹ distilled water) consisted of bacto saccharose, 30.0; sodium nitrate (NaNO₃), 3.0; dipotassium phosphate (K₂HPO₄), 1.0; magnesium sulfate (MgSO₄), 0.5; potassium chloride (KCl), 0.5; and ferrous sulfate (FeSO₄·7H₂O), 0.01. The flasks were incubated for 48 h at 30°C on a rotary shaker at 150 rpm in the dark.

ANALYTICAL METHODS AND DEGRADATION EXPERIMENTS

Isolation, detection, and identification of metabolite

To identify α -naphthol and metabolites, the fungus was cultivated in 250 ml Erlenmeyer flasks containing 100 ml of Czapek-Dox medium and inoculated with 2×10^8 conidia/ml medium. The α -naphthol (7.5 mg) was dissolved in 375 μ l of ethyl acetate, with Tween 80 (400 mg L⁻¹) added to one flask after 24 h of incubation in darkness. Five days after the addition of α -naphthol, the suspension was extracted three times, consecutively, with ethyl acetate. The organic extracts were combined after the separation of organic and water phases. Afterwards, the organic extracts were concentrated at 50°C by means of a rotary evaporator to about 5 ml [16]. Residues (5 ml) were examined for α -naphthol and acetate-extractable transformation products by GC and GC-MS. Abiotic controls (without microorganism) were always included.

Analysis of the sample with GC-MS

The samples were analyzed using a Hewlett-Packard 6890 with a DB-5 capillary column (30 m \times 0.25 mm; film thickness 0.25 μ m) was programmed as follows: 60°C for 5 min and 220°C at a rate of 4°C/min. The flow rate to helium as a carrier gas with (2 ml/min) MS was taken at 70 eV.

RESULTS

Biotransformation of α -naphthol by *A. niger*

In this experiment, the biotransformation of α -naphthol by *A. niger* grown on Czapek' s medium for only 5 days was performed. After incubation, Czapek' s culture was extracted (see analytical methods and degradation experiments section). The suspension was extracted with ethyl acetate three consecutive times and directly analyzed by GC-MS (fig. 1). In these analyses, various chemicals were obtained (tab. 1). The main products obtained in the bioconversion of *A. niger* of α -naphthol were 1-ethyl-2-methyl benzene (41%) and acetonephthone (7.43%), respectively.




Figure 1.
Mass spectrum of biotransformation of α -naphthol by *A. niger*

Table 1.

Mass spectral data, retention times, and proposed identities of metabolites derived from α -naphthol after 5 days of incubation

Proposed compound	Retention time (min)	Mass spectral ions, m/z (% of relative intensity)
1-Ethyl-2-methyl benzene	10.45	105(100), 120(53.58), 119(13.51), 77(10.63), 91(9.27)
Acetonaphthone	13.73	57.1(100), 43.1(72.28), 71.1(59.42), 85.1(35.96), 41.1(34.23)
1,4-Naphthoquinone	17.33	158(100), 102(45.43), 104(43.56), 130(33.49), 76(33.46)
4-Hydroxy-1-naphthyl sulfate	18.52	144(100), 115(80.27), 116(37.31), 145(10.85), 89(8.79)
2-Phenyl-1,2,3-tetrahydro-1-naphthol	19.40	105(100), 134(86.38), 77(31.34), 133(27.28), 120(23.89)




Figure 2.

Transformation of 1-naphthol by *A. niger*: 4-hydroxy-1-naphthylsulfate, 2-phenyl-1,2,3-tetrahydro-1-naphthol, 1,4-naphthoquinone and acetonaphthone were deduced from mass spectral data of 1-naphthol transformation products found in *A. niger* (tab. 1)

In a previous study of biotransformation of menthol by sporulated surface cultures of *A. niger* and *Penicillium* spp. the main bioconversion product obtained from menthol of *A. niger* was *cis-p*-menthan-7-ol, and the main products obtained by sporulated surface cultures of *Penicillium* spp. were limonene, *p*-cymene, and γ -terpinene [22]. Leuenberger reported that product yields could be effectively increased by solubilizing/emulsifying immiscible substrates [17]. However, careful selection of the nature and concentration of the solvent is necessary, because many miscible solvents are cytotoxic at lower concentrations [17]. Comparing the abovementioned investigation with the present study showed oxygenated monoterpenes to be the main compounds, with more monoterpenes yielded in the biotransformation. Figure 2 shows that transformation of α -naphthol by *A. niger* PTCC 5011 produces more sulphation products.

DISCUSSION

In previous research on biotransformation of α -naphthol by a strictly aquatic fungus studied the biotransformation of the environmental pollutant metabolite 1-naphthol has been studied. *H. lugdunensis* metabolized approximately 74% of 1-naphthol within 5 days [4]. The current research addressed the metabolism of α -naphthol by *A. niger* PTCC 5011, a coelomycete isolated in Ardebil, Iran, from industrial wastewater.

The study of microbial transformation of one monoterpene by sporulated surface cultures of *A. niger* and *Penicillium* sp. produced *cis-p*-menthan-7-ol from *A. niger*; the main products obtained were limonene, *p*-cymene, and γ -terpinene [11]. The two main products of microbial transformation of citral were similar to those obtained in former work. The main bioconversion products of (-)-menthol by *Mucor ramannianus* using the sporulated surface cultures method were *trans-p*-menthan-8-ol, *trans-menth-2-en-1-ol*, sabinane, *p*-menthane-3,8-diol, isomenthol, and 1,8-cineole [13].

The experimental work (the two latest articles) suggested that microbial transformation of monoterpenes with different genera: *Penicillium* and *Aspergillus* caused an oxidation reaction and resulted in a more stable product. Although, bioconversion by using *A. niger* PTCC 5011 showed that it was possible to obtain two or three main products of high percentage and selectivity.

CONCLUSION

In conclusion, reported literature and this research show bioconversion of α -naphthol by *A. niger* have been similarly effective for biotransformation. Oxidation and sulphation are two common characteristics in this investigation of bioconversion products.

ACKNOWLEDGMENTS

We thank Dr. Larijani, Department of Chemistry, Science & Research Campus, Islamic Azad University, Tehran, Iran, for performing the GC-MS measurements as well as Dr. Sheykhinejad and Mr. Parach, Department of Iranian Research Organization for Science & Technology, Tehran, Iran, for their support in this study.

REFERENCES

1. Akta N, Cicek H, Unal AT, Kibarer U, Kolankaya N, Tanyolaç A. Reaction kinetics for laccase-catalyzed polymerization of 1-naphthol. *Bioresource Tech* 2001; 80:29–36.
2. Allen CCR, Boyd DR, Larkin MJ, Reid AK, Sharma DN, Wilson K. Metabolism of naphthalene, 1-naphthol, indene, and indole by *Rhodococcus* spp. strain NCIMB 12038. *Appl and Env Microb* 1996; 63:151–155.
3. Anonymous A. *Sevin insecticide for abundant food and fibre*. Technical Information Bulletin, Union Carbide. 1974.
4. Augustin T, Schlosser D, Baumbach R, Schmidt J, Grancharov K, Krauss G, Krauss GJ. Biotransformation of α -naphthol by a strictly aquatic fungus. *Current Microb* 2005;52: 216–220.
5. Cerniglia CE, Althaus JA, Evans FE, Freeman JP, Mitchum RK, Yang SK. Stereochemistry and evidence for an arene oxide-NIH pathway in the fungal metabolism of naphthalene. *Chem Biol Inter* 1983; 44:119–132.
6. Cerniglia CE, Freeman JP, Evans FE. Evidence for an arene oxide-NIH shift pathway in the transformation of naphthalene to α -naphthol by *Bacillus cereus*. *Arch Microbiol* 1984; 138:283–286.
7. Cerniglia CE, Sutherland JB. Bioremediation of polycyclic aromatic hydrocarbons by ligninolytic and non-ligninolytic fungi. In: Gadd, G. M. (ed.) *British Mycological Society Symposium Series, Fungi in Bioremediation*. Cambridge 2001:136.
8. Crosby DG, Leitis E, Winterlin WL. Photo decomposition of carbamate insecticides. *J Agric Food Chem* 1965; 13:204-207.
9. Da Silva M, Esposito E, Moody JD, Canhos VP, Cerniglia CE. Metabolism of aromatic hydrocarbons by the filamentous fungus *Cylothryrium* spp. *Chemosphere* 2004; 57:934–952.
10. Doddamani HP, Ninnekar HZ. Biodegradation of Carbaryl by a *Micrococcus* species. *Current Microb* 2001; 43:69–73.
11. Esmaili A, Saad N, Safaiyan S, Rustaiyan A. Biotransformation of (-)-menthol by spores of *Mucor ramannianus* and study of the pathways involved. *Herba Pol* 2009; 56(2):51–57.
12. Esmaili A, Hoseiny Zarea A, Sharafian S, Safaiyan S, Rustaiyan A, Biotransformation of menthol by sporulated surface cultures of *Penicillium* spp. and study of the pathways involved. *Herba Pol* 2009; 55(1):78–83.
13. Esmaili A, Sharafian S, Safaiyan S, Rezazadeh S, Rustaiyan A. Biotransformation of one monoterpene by sporulated surface cultures of *Aspergillusniger* and *Penicillium* spp. *Nat Prod Res* 2009; 23:1058–1061.
14. Hasegawa Y, Okamoto T, Obata H, Tokuyama T. Utilization of aromatic compounds by *Trichosporon cutaneum*. *KUY* 1990; 69:122–126.
15. Hidalgo C, Sancho JV, Roig-Navarro A, Hernandez F. Rapid determination of carbaryl and α -naphthol at ppt levels in environmental water samples by automated on-line SPE-LC-DAD-FD. *Chromatographia* 1998; 47:596–600.
16. Joshi AA, Locke BRP, Finney WC. Formation of hydroxyl radicals, hydrogen peroxide and aqueous electrons by pulsed streamer corona discharge in aqueous solution. *J Hazard Mater* 1995; 41:3–30.
17. Leuenberger HGW. Methodology. In: Kieslich, K. (ed.). *Biotransformations*, vol. 6A. Weinheim 1984:5.
18. Mastrangelo G, Fadda E, Marzia V. Polycyclic aromatic hydrocarbons and cancer in man. *Environ Health Perspect* 1997; 104:1166–1170.

19. Muncnerova D, Augustin J. Fungal metabolism and detoxification of polycyclic aromatic hydrocarbons: a review. *Bioresour Technol* 1994; 48:97–106.
20. Preuss R, Angerer J, Drexler H. Naphthalene—an environmental and occupational toxicant. *International Archives of Occupat and Env Health* 2003; 76:556–576.
21. Samanta SK, Chakraborti AK, Jain RK. Degradation of phenanthrene by different bacteria: evidence for novel transformation sequences involving the formation of 1-naphthol. *Appl Microbiol Biotechnol* 1999; 53:98–107.
22. Sikka HC, Miyazaki S, Lynch RS. Degradation of Carbaryl and α -naphthol by marine microorganisms. *Bulletin of Env Contam & Toxicol* 1975; 13(6):666–672.
23. Zang S, Lian B. Synergistic degradation of 2-naphthol by *Fusarium proliferatum* and *Bacillus subtilis* in wastewater. *J Hazard Mat* 2008; 166:33–38.
24. Esmaeili A, Hashemi E. Biotransformation of myrcene by *Pseudomonas aeruginosa*. *Chem Central J* 2011; 5:26–34.
25. Esmaeili A, Hashemi E, Safaiyan SH, Rustaiyan A. Biotransformation of myrcene by *Pseudomonas putida* PTCC 1694. *Herba Pol* 2011; 57(1):51–58.
26. Esmaeili A, Hashemi E, Safaiyan SH, Rustaiyan A. Biotransformation of germacranolide from *Onopordon leptolepis* by *Aspergillus niger*. *Pak J Pharm Sci* 2012; 25(1):155-159.
27. Esmaeili A, Khodadadi A. Antioxidant activity of a solution of thymol in ethanol. *Zahedan Uniof Med Sci and Health Ser* 2012; 14(7):14-18.
28. Esmaeili A, Khodadadi A, Safaiyan S. Biotransformation of thymol by *Aspergillus niger*. *Chem Nat Comp* 2012; 47(6):966-968.
29. Esmaeili A, Saremnia B, Koohian A, Rezazadeh S. Mechanism of nanocapsules of *Matricaria recutita* extract formation by the emulsion-diffusion process. *Superlattices and Microstructures* 2011; 50:340–349.
30. Esmaeili A, Saremnia B. Preparation of extract-loaded nanocapsules from *Onopordon leptolepis* DC. *Ind Crops and Prod* 2012; 37:259–263.
31. Esmaeili A, Kalantari M. Bioremoval of an azo textile dye, Reactive Red 198, by *Aspergillus flavus*. *World J Microbiol Biotechnol* 2012; 28:1125–1131.

TRANSFORMACJA MIKROBIOLOGICZNA α -NAFTOLU PRZEZ *ASPERGILLUS NIGER* – PTCC 5011AKBAR ESMAEILI^{1,*}, SHAHROUZ FAZELI¹, NASRIN MOAZAMI²

¹Department of Chemical Engineering,
North Tehran Branch
Islamic Azad University
P.O. Box 19585-936
Tehran, Iran

²Department of Biotechnology
Iranian Research Organization for Science and Technology
Teheran, Iran

Streszczenie

Badano przetwarzanie α -naftolu za pomocą *Aspergillus niger* PTCC 5011, grzyba wyizolowanego ze ścieków przemysłowych pochodzących z Ardebil w Iranie. Kropidlak czarny metabolizował około 80% α -naftolu w ciągu pięciu dni. Identyfikacja i ocena jakościowa produktów degradacji za pomocą GC/MS wykazała, że około 41% związku macierzystego zostało przetworzone na 1-etyl-2-metylobenzen, 7,43% na acetonafton, 5,55% zostało przekształcone w 4-hydrokso-1-naftylosulfat, 3% w naftochinon i około 6,68% w 2-fenyl-1,2,3-tetrahydro-1-naftol. Takie wyniki mogą pomóc znaleźć zastosowanie *A. niger* do regulacji wpływu zanieczyszczeń typu tłuszczowego, które istnieją w ekosystemach.

Słowa kluczowe: biotransformacja, biokonwersja, *Aspergillus niger*, α -naftol, grzyby